

Download File PDF Cpr Questions And Answers For Exam Documenter

#Jenny



Finally I get this ebook, thanks for all these I can get now!

#Rio



Cool! I'am really happy

#Markus Jensen



I did not think that this would work, my best friend showed me this website, and it does! I get my most wanted eBook

#Hun Tsu



wtf this great ebook for free?!

#Che Salsa



My friends are so mad that they do not know how I have all the high quality ebook which they do not!

#Diego Butler



so many fake sites. this is the first one which worked! Many thanks

[Download PDF version of :](#)
Cpr Questions And Answers For Exam Documenter

- CPR and AED Test**
1. Which action is more likely to cause the rescuer?
 - a. Does not make good contact between face and mouth
 - b. Does breathe too quickly with too much force
 - c. Does not breathe fast enough
 - d. Does breathe too fast
 2. The compression-to-ventilation ratio for 2-rescuer adult CPR is:
 - a. 30:2
 - b. 15:2
 - c. 30:1
 - d. 15:1
 3. If a victim of a large body injury (fracture) becomes unresponsive, the rescuer should send someone to activate the emergency response system and immediately:
 - a. Call the medical director
 - b. Perform advanced airway
 - c. Perform head-tilt/chin-lift
 - d. Start CPR (beginning with compressions)
 4. When performing CPR with the correct compression and ventilation rate for 2-rescuer CPR in the presence of an advanced airway:
 - a. Compress at a rate of at least 100 per minute, 1 breath every 6 to 8 seconds
 - b. Compress at a rate of at least 100 per minute, 2 breaths every 5 to 10 seconds
 - c. Compress at a rate of at least 100 per minute, 1 breath every 10 to 12 seconds
 - d. Compress at a rate of at least 100 per minute, 1 to 2 breaths every 6 to 8 seconds
 5. The compression-to-ventilation ratio for 2-rescuer adult CPR is:
 - a. 30:2
 - b. 15:2
 - c. 30:1
 - d. 15:1
 6. The 2020 AHA guideline for CPR recommends BLS sequence of steps are:
 - a. Airway, Breathing, Check Pulse
 - b. Airway, Breathing, Chest Compressions
 - c. Check compression, Airway, Breathing
 - d. None of the Above
 7. When a child has a heart rate greater than 100 per minute and a pulse that is not breathing effectively, the rescuer should:
 - a. Continue the AED on the child and analyze
 - b. Give breaths and chest compressions
 - c. Give breaths without chest compressions
 - d. Give chest compressions with breaths
 8. When should a rescuer attempt to palpate the brachial pulse in an infant?
 - a. Inside the upper arm, between the elbow and the shoulder
 - b. On the middle side of the upper leg, near the groin
 - c. On the side of the lower arm, near the wrist
 - d. On the side of the neck, near the trachea
 9. If an AED is used for a child, the rescuer should:
 - a. Adult pads may be used, but they should be cut in half before application.
 - b. Adult pads may be used if pediatric pads are not available.